

Project: **1178**

Project title: **Analysis of representation of wind fields in the models participating in DYAMOND summer, DYAMOND winter and nextGEMS**

Principal investigator: **Roland Schrödner**

Report period: **2023-05-01 to 2025-04-30**

The project was outlined for two main tasks: i) statistical analysis of the wind fields simulated by the models participating in DYAMOND summer, DYAMOND winter as well as nextGEMS (<https://easy.gems.dkrz.de>) and ii) investigating the effects on modelled global mineral dust and sea spray emissions. As an additional project component, not present in last year's proposal, iii) an analysis of ice nucleating particles (INP) and underlying aerosol conditions from global aerosol simulations (ECHAM, EC-Earth) were conducted. The project is therefore mainly a data analysis project, which requires storage capacities as the simulation data of several TB cannot easily be transferred to other machines. The project did not require and use large computational resources. The project tasks were and are mainly conducted by student helpers and are therefore not continuously worked on. Further, this lead to an adaptation of the project tasks to include analysis of existing other simulations for aerosols and INP.

In the reporting period, in particular analysis for task i) was conducted, which was continued for both DYAMOND scenarios.

In addition to the originally planned analyses, the simulations with the global atmospheric model ECHAM (done under resources of a different project) and EC-Earth (available from a recent publication) were further post-processed and these data saved for use in this project. During the project duration, most of the planned storage resources were used temporarily. Computational resources were not used and less will be applied for in the upcoming allocation period. The already analyzed data that is saved under the project's resources is still of use and it is aimed to further analyze it. Therefore, similar storage resources will be applied for. The content of the request for the upcoming period will be largely unchanged.

Results of statistical analysis of DYAMOND model's wind fields (covering the whole project period beyond the last year)

For the analysis, the vertical wind fields were translated into histograms showing frequencies of occurrence with 0.1 m s^{-1} resolution for all models and available horizontal resolutions. Apart from the global field (3D + time), also subdomains (lowest 500 m vs full troposphere, Central Europe) were analyzed similarly. In addition, such histograms were created for the high-resolution regional simulations performed with the ICON-LEM model over Germany during the HD(CP)2 project, which are also hosted at DKRZ. The different model data were compared to observations of vertical velocity derived from doppler lidars and windprofilers. The observational datasets comprise several years and is representative for Central German Lowlands (observations situated at Leipzig and Lindenberg, Germany). From the observational datasets comparable histograms were created.




Figure 1: PDF of vertical velocity (x-axis in m s^{-1}) of different resolutions of the global ICON runs from DYAMOND and regional ICON LEM runs from HD(CP)2, and comparison against lidar and windprofiler observations in Leipzig.

The comparison revealed a large spread between the different high-resolution models in terms of the frequency of occurrence of strong vertical wind speeds ($\pm 5 \text{ m s}^{-1}$, factor of 2-3 for updrafts and

factor 10-100 for downdrafts). As expected also the model resolution has a large impact on the occurrence rate of vertical velocities due to the more smoothed wind fields. In comparison to the observed vertical velocity frequencies at the two locations in Germany, it could be seen that none of the global high resolution models reaches similar high frequencies in the observable range with sufficient statistics between $\pm 0.5 - 7 \text{ m s}^{-1}$ (and opposite overestimating low vertical velocity frequencies). Naturally, the available horizontal resolutions of 2.5 km and more miss the observed small scale structures. Observation and model can be brought closer together, if the observations are averaged to time intervals of a few minutes. Furthermore, the vertical velocities frequencies derived from the 156 m ICON-LEM simulations show a good agreement with the observed frequencies. However, the interpretation needs some caution since the selection of simulated cases with ICON-LEM during HD(CP)2 is not necessarily representative to the observation period of several years. However, the occurrence of high vertical velocities is mostly driven by convective situations, which was a key target for the HD(CP)2 project.

A publication on this analysis is still in preparation, but on hold until nextGEMS simulations have been analyzed accordingly. The analysis was refined in terms of vertical extent of the conducted histograms in order to match observed and modeled vertical velocities better.


In addition to vertical velocities, horizontal wind fields of some of the DYAMOND summer simulations were extracted and histograms were produced. The extracted fields were applied to a desert dust emission algorithm typically used in atmospheric chemistry transport models.

Further analysis in the next allocation period aims at taking into account as well the DYAMOND winter and nextGEMS runs.

Results of analysis of aerosol simulation data for diagnosing INP

In the reporting period, global aerosol simulation data of the global models ECHAM and EC-Earth were used to diagnose spatiotemporally varying fields of INP concentrations. For this purpose, INP parameterizations were applied to certain aerosol data proxies (mineral dust, sea spray aerosol). The resulting INP-temperature-spectra were compared to available observational data. The utilized simulation data sets comprise 1 year with hourly output (EC-Earth) and 30 years with 6-hourly output of aerosol concentrations with a horizontal resolution of roughly $2^\circ \times 3^\circ$. The analyzed INP data set was used in a recent publication (Hartmann, Schrödner, Zeppenfeld et al., 2025). Further the cloud and temperature fields of the 30-year ECHAM runs were investigated for the likelihood of the occurrence of secondary ice production in clouds, which was used in a Bachelor Thesis.

The analysis of these long-term datasets shall be continued by taking into account further INP



observations and other parameterizations, e.g., for bioaerosol. Therefore the title of the upcoming request was adapted.

Figure 2: Modelled vs. observed INP concentrations at different Southern Ocean sites between -15 and -20°C . The grey point cloud shows the INP resulting from modeled mineral dust and the coloured points for in addition to mineral dust taking into account marine polysaccharides present in sea spray aerosol. The numbers above the plot show the agreement within a factor of 2 and a factor of 10, hence, depicting the improvement by taking into account the newly implemented polysaccharidic INP.

Hartmann, S., Schrödner, R., Hartmann, M., van Pinxteren, M., Fomba, W., Stratmann, F., Herrmann, H., Pöhlker, M., Hassett, B. T., and Zeppenfeld, S. (2025): Marine polysaccharides - Important Constituents of Ice Nucleating Particles of Marine Origin, *Env. Sci. & Tech.*, 59 (10), 5098-5108, <https://doi.org/10.1021/acs.est.4c08014>.